

Entropy radiated by a braneworld black hole

Aurélien Barrau,^{1,*} Julien Grain,^{2,†} and Carole Weydert^{1,‡}

¹*Laboratoire de Physique Subatomique et de Cosmologie,
UJF, CNRS/IN2P3, INPG*

53 avenue des Martyrs, 38026 Grenoble cedex, France

²*AstroParticule et Cosmologie, Université Paris 7, CNRS/IN2P3
10, rue Alice Domon et Léonie Duquet, 75205 Paris cedex 13, France*

(Dated: February 2, 2008)

The concept of black hole entropy is one of the most important enigmas of theoretical physics. It relates thermodynamics to gravity and allows substantial hints toward a quantum theory of gravitation. Although Bekenstein conjecture –assuming the black hole entropy to be a measure of the number of precollapse configurations– has proved to be extremely fruitful, a direct and conclusive proof is still missing. This article computes accurately the entropy evaporated by black holes in $(4 + n)$ dimensions taking into account the exact greybody factors. This is a key process to constrain and understand the entropy of black holes as the final state is unambiguously defined. Those results allow to generalize Zurek’s important argument, in favor of the Bekenstein conjecture, to multi-dimensional scenarios.

PACS numbers: 04.70.Dy, 04.62.+v, 04.70.-s, 04.50.+h

This article aims at generalizing to an arbitrary number of dimensions the key argument on the entropy of black holes made by Zurek [1], improved by Page [2] and reconsidered by Schumacher [3]. This is a mandatory step to consider black holes in braneworld scenarios. As it requires quite sophisticated numerical techniques to compute the greybody factors, this point could not have been easily made earlier.

Bekenstein [4] proposed that black holes possess an entropy proportional to their surface area, which can be statistically interpreted as a measure of the number of precollapse configurations leading to a given state. It can easily be shown that the only possible power-law function of the area A allowed for the entropy is linear (see, *e.g.* [5], for an introductory review). The constant of proportionality can be fixed using the Hawking law [6], leading to $S = A/4$ in Planck units. Attempts to understand this simple formula from more fundamental points of view are legion. Important historical points were made by Bombelli *et al.* [7] and Srednicki [8] to relate the entropy of black holes to entanglement entropy arising from the tracing out of those degrees of freedom localized beyond the horizon. Thorne and Zurek [9] and ’t Hooft [10] considered the thermal radiative “atmosphere” of black holes. More recently, the entropy has been linked with the degrees of freedom of strings associated with the hole [11, 12] or with the quantum gravity degrees of freedom of the horizon, in the context of conformal field theory [13, 14] or in loop quantum gravity [15]. More heuristic schemes were also developed, *e.g.* in [16]. Basically speaking, recovering the Bekenstein black hole entropy is one of the most important successes of the main tentative theories of quantum gravity. This happens quite naturally in string theory (at least for *some* extremal black holes – see [17, 18] for reviews) and in loop quantum gravity (at least when setting correctly the free Immirzi parameter – see [19]).

However, a direct proof of Bekenstein’s interpretation of the entropy (relating its meaning with the outer world) is still missing. Zurek’s idea to go further in this direction was to consider the black hole evaporation and to quantify the ratio of the evaporated entropy to the entropy lost by the black hole [1]. This is a fundamental process to this purpose because the final state is well determined and its entropy can be accurately estimated. The key ingredient for this computation –namely the greybody factors– were derived by Page [20] (following Starobinsky [21]), thanks to exhaustive numerical computations, allowing to solve this problem exactly in 4 dimensions [2, 20]. However, quite a lot of modern fundamental models of theoretical physics –beginning of course with string theory– involve extra-dimensions. We now proceed to the investigation of this problem in the Arkani-Hamed-Dimopoulos-Dvali (ADD) scenario [22] based on large extra-dimensions which accounts, among other problems, for the huge hierarchy between the different scales of physics. The keypoint is to allow only gravitons to propagate in the extra dimensions while standard model fields are localized on the brane.

*Electronic address: Aurelien.Barrau@cern.ch

†Electronic address: grain@apc.univ-paris7.fr

‡Electronic address: weydert@lpsc.in2p3.fr

The metric for a $(4 + n)$ -dimensional black holes reads as

$$ds^2 = -h(r)dt^2 + h(r)^{-1}dr^2 + r^2 d\Omega_{n+2}^2,$$

where $d\Omega_{n+2}$ is the solid angle element and $h(r) = 1 - (r_H/r)^{n+1}$, with r_H the horizon radius. The temperature T of such black holes is given by

$$T = \frac{1}{4\pi} \frac{dh}{dr} \Big|_{r_H} = \frac{n+1}{4\pi r_H},$$

the radius being related with the mass M of the hole through

$$r_H = \frac{1}{\sqrt{\pi} M_*} \left(\frac{M}{M_*} \right)^{\frac{1}{n+1}} \left(\frac{8}{n+2} \right)^{\frac{1}{n+1}} \left[\Gamma \left(\frac{n+3}{2} \right) \right]^{\frac{1}{n+1}},$$

where M_* is the fundamental $(4 + n)$ -dimensional Planck scale, while the area is given by

$$A = r_H^{n+2} (2\pi)^{\frac{n+1}{2}} \left[\Gamma \left(\frac{n+3}{2} \right) \right]^{-1}.$$

As a first approximation, for particles confined on the brane, the radiation emitted in a short time interval dt carries an energy $dE = \Sigma \alpha T^4 dt$ and an entropy $dS_{rad} \approx \frac{4}{3} \Sigma \alpha T^3 dt$ where Σ is an effective cross section and α is the Stefan constant. This allows to compute the variation of the black-hole entropy:

$$\begin{aligned} dS_{BH} &= \frac{dA}{4} M_*^{n+2} = \frac{M_*^{n+2}}{4} [A(M) - A(M - dE)] \\ &\approx f(n) \frac{n+2}{n+1} M^{\frac{1}{n+1}} dE \end{aligned}$$

with

$$f(n) = \frac{\sqrt{\pi}}{2} \frac{1}{M_*^{\frac{n+2}{n+1}}} \left(\frac{8}{n+2} \right)^{\frac{n+2}{n+1}} \left[\Gamma \left(\frac{n+3}{2} \right) \right]^{\frac{1}{n+1}}.$$

The ratio R of the entropy lost by the black hole to the entropy gained by the radiation can be estimated:

$$R = \frac{dS_{rad}}{dS_{BH}} = \frac{4}{3} \frac{n+1}{n+2} \frac{1}{f(n)T} M^{-\frac{1}{n+1}} = \frac{4}{3}.$$

This is exactly the same result as obtained by Zurek in 4 dimensions. As one could expect, the effect of the modification of the area of the black hole due to the extra-dimensions is totally compensated by the modification of its temperature. However, it will be demonstrated that a more rigorous treatment of this problem radically modifies this conclusion. This is due both to the emission of gravitons in the bulk and to the footprint of the extra-dimensions on the greybody factors, even for brane fields.

The exact semi-classical flux spectrum of a higher-dimensional black hole is given by:

$$\frac{dN^{(s)}(\omega)}{dt} = \sum_{\ell, m} \sigma_n^{(s)}(\omega) \frac{1}{\exp(\frac{\omega}{T}) - (-1)^{2s}} \frac{d^{n+3}k}{(2\pi)^{n+3}}$$

where the cross section σ accounts for the couplings between the black hole and quantum fields with spin s . Using the "macroscopic" thermodynamical approach of Zurek, one can evaluate R for standard model fields through

$$R = \frac{dS_{rad}}{dS_{BH}} = \frac{\int_0^\infty \sigma \left[\frac{x}{e^x - 1} - \ln(1 - e^{-x}) \right] x^2 dx}{\int_0^\infty \sigma \frac{x^3}{e^x - 1} dx}.$$

The results are given, as a function of the number of extra-dimensions, in Fig. 1. A weak dependance upon n can be noticed, together with a good agreement with the black-body approximation.




FIG. 1: "Macroscopic" ratio of the radiated entropy to the variation of the hole entropy, as a function of the number of extra-dimensions for fermions and gauge bosons confined on the 4-brane.

Although more meaningful than the first approximation, this approach still misses the fundamental point, *i.e.* an accurate account for the greybody factors when the entropy is correctly expressed with the density matrix as $tr(-\rho \ln \rho)$. For a non-rotating, uncharged, black hole the radiation is emitted with a density matrix in each mode

$$\rho_{kk'} = \delta_{kk'} \frac{|A_{\ell,s}|^{2k} (e^{\frac{\omega}{T}} - (-1)^{2s})}{(e^{\frac{\omega}{T}} - (-1)^{2s} + (-1)^{2s} |A_{\ell,s}|^2)^{k+(-1)^{2s}}}$$

where $|A_{\ell,s}|^2$ is the transmission coefficient of the black hole to the considered particle. This allows to write

$$\frac{dS_{BH}}{dt} = -\frac{1}{T} \frac{dM}{dt} = -\frac{1}{T} \sum_{\ell,m,s} \int \frac{d\omega}{2\pi} \omega \underbrace{\frac{|A_{\ell,s}|^2}{e^{\omega/T} - (-1)^{2s}}}_{\langle N_{\omega,\ell,m} \rangle},$$

where the summation over the angular momentum ℓ and its projection m must be carried out after performing the integrals, and:

$$\frac{dS_{rad}}{dt} = \sum_{\ell,m,s} \int \frac{d\omega}{2\pi} \left[\underbrace{\frac{|A_{\ell,s}|^2}{e^{\omega/T} - (-1)^{2s}} \ln \left(\frac{e^{\omega/T} - (-1)^{2s}}{|A_{\ell,s}|^2} + (-1)^{2s} \right) + (-1)^{2s} \ln \left(1 + (-1)^{2s} \frac{|A_{\ell,s}|^2}{e^{\omega/T} - (-1)^{2s}} \right)}_{\langle S_{\omega,\ell,m} \rangle} \right].$$

The coefficients $A_{\ell,s}$ account both for the intricate metric structure around the black hole and for the centrifugal barrier, possibly leading to a back-scattering of the emitted particle [23]. They can be determined by solving the equations of motion with appropriate boundary conditions at the black hole's horizon and at spatial infinity. For particles confined on the brane –with the usual ansatz $\Phi_s(t, r, \Omega) = e^{-i\omega t} Y_{m,s}^\ell(\Omega) R_s(r)$ to split the temporal, angular and radial parts of the field (where $Y_{m,s}^\ell$ are the spin-weighted spherical harmonics)– the radial function $R_s(r)$ obeys the master equation:

$$\left[\Delta^s \frac{d}{dr} \left(\Delta^{1-s} \frac{d}{dr} \right) + \left(\frac{\omega^2 r^2}{h} + 2is\omega r - \frac{is\omega r^2}{h} \frac{dh}{dr} - \lambda \right) \right] P_s = 0$$

where $\Delta = r^2 h(r)$, $P_s(r) = \Delta^s R_s(r)$ and $\lambda = j(j+1) - s(s-1)$. This is equivalent, in the tortoise coordinate system, to a Shrödinger equation with a potential depending on the metric and its derivatives, on the number of

extra-dimensions and on the spin of the field. The keypoint is that the derived greybody factors keep a clear footprint of the dimensionality of space, even for those particles living on the 4-brane. This is the main reason why the entropy ratios are strongly modified by the existence of extra-dimensions. The case of gravitons –propagating in the bulk– is more subtle, requiring to investigate the three kind (scalar, vector, tensor) of gravitational perturbations evolving in a higher dimensional space (see, *e.g.*, Ref. [24, 25, 26]). Although analytical expressions can be obtained in the infrared and ultra-violet limits for all the aforementioned cases, the most interesting regime, where $\omega r_H \sim 1$, requires dedicated numerical investigations for each kind of particle. We have solved the appropriate field equations with an extensive simulation (see, *e.g.*, [27] for a description) which, basically, proceeds in three steps: for each value of the energy ω and of the angular momentum ℓ , the radial part of the equations is numerically solved, then the amplitudes of the outgoing and ingoing modes at spatial infinity are estimated by fitting the asymptotic analytical solutions and, finally, the resulting transmission coefficient is used to compute the contribution to the ℓ -th multipole. Some examples for 0, 1 and 7 extra-dimensions are given in Fig. 2.




FIG. 2: Cross-sections as a function of the energy of the particle multiplied by the horizon radius for $n=0$, $n=1$ and $n=7$. Upper-left: scalars on the brane, upper right: fermions on the brane, lower left: gauge bosons on the brane, lower right: gravitons in the bulk. For this latter plot the three curves correspond (respectively from top to bottom at $\omega r_h = 2$) to the tensor, vector and scalar contributions.

Using those transmission coefficients, it is possible to compute exactly the entropy lost by the black hole and gained by the radiation. Results are given in Tab. I, as a function of the number of degrees of freedom, and constitute the first computation of the entropy radiated by a braneworld black hole. This is the key result of this paper. Remarkably, this process allows to relate the –speculative– entropy of the black hole in $(4 + n)$ dimensions with the well known entropy of a multidimensional blackbody radiation. Throughout this paper the emitted particles are assumed to be massless. Although taking into account the mass will inevitably modify the greybody factors [28], this will not change our conclusions as the point remains valid for most black hole temperatures. When the black hole is very massive

($T \ll \mu_{min}$ where μ_{min} is the mass of the lightest particle) only massless fields will be radiated and when the black hole is very light ($T \gg \mu_{max}$ where μ_{max} is the mass of the heaviest particle) ultra-relativistic quanta will be radiated. Important modifications of the transmission coefficients are therefore only expected when $T \in [\mu_{min}, \mu_{max}]$.




FIG. 3: Exact computation of the ratio of the radiated entropy to the variation of the black hole entropy, as function of the number of extra-dimensions for scalars, fermions and gauge bosons confined on the 4-brane and gravitons propagating in the bulk.

The ratios $R = dS_{rad}/dS_{BH}$ are plotted in Fig. 3. It should be noticed that, contrarily to what happens in Zurek's approach, nothing here mathematically prevents R from being arbitrarily large or small. The fact that the values computed are very close to one is a strong point in favor of the conjecture of Bekenstein. Furthermore, it is still possible –at least at the *gedankenexperiment* level– to consider a black hole in thermal equilibrium with a surrounding heat bath. The condition of equilibrium simply requires the energy associated with the radiation to be much smaller than the mass of the hole [29]. This straightforwardly generalizes to an arbitrary number of dimensions. In this situation, adiabatically changing the volume of the container so that the entropy production ratio remains arbitrarily close to one allows to reinforce the validity of the generalized second law of thermodynamics.

Finally, one could also play the game to consider a black hole emitting all the fields in all the extra-dimensions. An elementary thermodynamical consideration allows to prove that, at the first order of approximation, $R = \frac{n+4}{n+3}$. It means that, as the number of dimensions increases, the evaporation process becomes more and more isentropic. Or, the other way round, that the subtle correlations assumed in 4 dimensions to account for the "artificial" entropy increase during the evaporation process become less and less important. This would mean that the coarse-grained entropy becomes equivalent to the fine-grained entropy, therefore solving the unitarity problem. A more refined numerical study shows that this estimate of R is correct at the percent level.

n	dS_{rad}/dt ($10^{-3} r_H^{-1}$)	dS_{BH}/dt ($10^{-3} r_H^{-1}$)
0	$(6.948 n_0 + 3.368 n_{1/2} + 1.269 n_1 + 0.260 n_2)$	$(3.740 n_0 + 2.060 n_{1/2} + 0.846 n_1 + 0.193 n_2)$
1	$(30.32 n_0 + 24.39 n_{1/2} + 17.87 n_1 + 13.10 n_2)$	$(16.77 n_0 + 14.57 n_{1/2} + 11.48 n_1 + 9.712 n_2)$
2	$(79.12 n_0 + 68.35 n_{1/2} + 64.19 n_1 + 88.75 n_2)$	$(45.71 n_0 + 41.44 n_{1/2} + 40.80 n_1 + 66.33 n_2)$
3	$(160.7 n_0 + 139.8 n_{1/2} + 147.5 n_1 + 326.3 n_2)$	$(97.57 n_0 + 88.34 n_{1/2} + 94.27 n_1 + 246.4 n_2)$
4	$(280.8 n_0 + 245.4 n_{1/2} + 273.1 n_1 + 988.2 n_2)$	$(179.3 n_0 + 164.4 n_{1/2} + 176.9 n_1 + 760.8 n_2)$
5	$(445.4 n_0 + 394.2 n_{1/2} + 446.1 n_1 + 2920 n_2)$	$(297.2 n_0 + 281.3 n_{1/2} + 294.1 n_1 + 2305 n_2)$
6	$(660.7 n_0 + 595.6 n_{1/2} + 671.0 n_1 + 8881 n_2)$	$(457.1 n_0 + 447.4 n_{1/2} + 449.9 n_1 + 7183 n_2)$
7	$(932.3 n_0 + 846.0 n_{1/2} + 950.7 n_1 + 28386 n_2)$	$(661.3 n_0 + 646.5 n_{1/2} + 645.0 n_1 + 23466 n_2)$

TABLE I: Rates of entropy variation for the radiation and for the black hole, in units of $10^{-3} r_H^{-1}$ as a function of the number of extra-dimensions n and of the scalar (n_0), fermionic ($n_{1/2}$), bosonic (n_1) and gravitational (n_2) degrees of freedom. Gravitons are assumed to propagate in the bulk whereas the other fields propagate in the brane.

This article provides the first computation of the entropy evaporated by a braneworld black hole. The results are given at three different levels of approximation, the latter taking into account the complete numerical evaluation of the greybody factors. This relates entropies of totally different conceptual origins, one being gravitational –associated with the area of the black hole– and the other being statistical –associated with the emitted radiation. This is a new step toward a full understanding of the entropy of black holes in multi-dimensional scenarios required by most particle physics and cosmology unification models. Remarkably, it does not rely on an assumed fundamental theory (*e.g.* strings or loops) but just on the well established semi-classical evaporation of black holes. This favours the conjecture of Bekenstein in $(4+n)$ -dimensions and, most importantly, allows exact computations of the entropy ratios.

-
- [1] W.H. Zurek, Phys. Rev. Lett. 49 (1982) 1683
 - [2] D.N. Page, Phys. Rev. Lett. 50 (1983) 1013
 - [3] B.W. Schumacher, Phys. Rev. Lett. 54 (1985) 2643
 - [4] J.D. Bekenstein, Phys. Rev. D 7 (1973) 2333
 - [5] J.D. Bekenstein, Contemp. Phys. 45 (2003) 31
 - [6] S.W. Hawking, Commun. Math. Phys. 43 (1975) 199
 - [7] L. Bombelli, R.K. Koul, J. Lee & R.D. Sorkin, Phys. Rev. D 34 (1986) 373
 - [8] M. Srednicki, Phys. Rev. Lett. 71 (1993) 666
 - [9] W.H. Zurek & K.S. Thorne, Phys. Rev. Lett. 54 (1985) 2171
 - [10] G. 't Hooft, Nucl. Phys. B 256 (1985) 727
 - [11] A. Strominger & C. Wafa, Phys. Lett. B 379 (1996) 99
 - [12] J. Maldacena, A. Strominger & E. Witten J. High Energy Phys. 9712 (1997) 002
 - [13] S. Carlip, Class. Quantum Grav. 16 (1999) 3327
 - [14] S.N. Solodukhin, Phys. Lett. B 454 (1999) 213
 - [15] A. Ashtekar, J. Baez, A. Corichi & K. Krasnov, Phys. Rev. Lett. 80 (1998) 904
 - [16] J.D. Bekenstein & V.F. Mukhanov, Phys. Lett. B 360 (1995) 7
 - [17] T. Damour, hep-th/0401160
 - [18] J.R. David, G. Mandal & S.R. Wadia, Phys. Rep. 369 (2002) 549
 - [19] A. Ashtekar, J.C. Baez & K. Krasnov, Adv. Theor. Math. Phys. 4 (2000) 1
 - [20] D.N. Page, Phys. Rev. D 13 (1976) 198
 - [21] A.A. Starobinsky, Sov. Phys. - JETP 37 (1973) 28
 - [22] N. Arkani-Hamed, S. Dimopoulos & G.R. Dvali, Phys. Lett. B 429 (1998) 263
 - [23] P. Kanti, Int. J. Mod. Phys. A 19 (2004) 4899
 - [24] H. Kodama & A. Ishibashi, Prog. Theor. Phys. 110 (2003) 701
 - [25] S. Creek, O. Efthimiou, P. Kanti & K. Tamvakis, Phys. Lett. B 635 (2006) 39
 - [26] V. Cardoso, M. Cavaglia & L. Gualtieri, JHEP 0602 (2006) 021
 - [27] J. Grain, A. Barrau & P. Kanti, Phys. Rev. D 72 (2005) 104016
 - [28] J. Grain & A. Barrau, arXiv:hep-th/0701265v1, in press for Class. Quantum Grav.
 - [29] S.W. Hawking, Phys. Rev. D 13 (1976) 191